arXiv:hep-ph/0201159v3 4 Jun 2002

ISU-HET-02-1
hep-ph/0201159
June, 2002

Probing Intrinsic Charm with Semileptonic B Decays

F. Gabbiani, Jianwei Qiu, and G. Valencia

Department of Physics and Astronomy, Iowa State University, Ames, Iowa 50011

Abstract

We discuss semileptonic B decays of the form $B \to J/\Psi\ e\nu\ X$ as possible probes of intrinsic charm. We calculate the leading order perturbative contribution to the process $B \to J/\Psi\ e^- \overline{\nu}_e\ X$ and find it to be unobservably small, with a branching ratio $\sim 10^{-10}$. We propose a modified spectator model to estimate the intrinsic charm contribution and find that it can be significantly larger, with a branching ratio for $B \to (c\bar{c})\ e^- \overline{\nu}_e\ X$ as large as 5×10^{-7}. We show that the process could be observed at these levels by CDF assuming a Run II integrated luminosity of 15 fb^{-1}, making this a useful reaction to probe the idea of intrinsic charm.

1 Introduction

Interesting possibilities open up if we consider wavefunctions of hadronic bound states that contain light-cone Fock states of arbitrarily high particle number. The higher light-cone Fock components arise as quantum fluctuations suppressed by M^2, where M is the mass of the fluctuation. The $c\bar{c}$ component of hadronic Fock states is referred to as intrinsic charm (IC) [1].

The possibility of detecting signatures of IC in B-meson physics has been recently discussed in Ref. [2]. The authors propose IC as the explanation for a "slow" (low momentum) bump in the inclusive $B \rightarrow J/\Psi\, X$ spectrum and the softness of the J/Ψ spectrum in $\Upsilon \rightarrow J/\Psi\, X$ decays, contrary to expectations from the color octet mechanism. IC appearing in higher light-cone Fock components of the B-meson wavefunction can manifest itself in two possible ways. It can operate virtually, in a mediation rôle, and affect decay processes by providing additional channels for the weak interactions. In this way one may enhance some CKM suppressed B decays. This has been discussed in Ref. [3] using a phenomenological mixing angles approach, and more recently in Ref. [4]. Another instance of this mediation is given by the hypothesis that the IC component of ρ mesons is able to explain the "$\rho\pi$ puzzle" [5]. IC can also manifest itself in processes in which the (intrinsic) $c\bar{c}$ produces charmed hadrons in the final state. An example of this is discussed in Ref. [6] to account for charm production in deep inelastic scattering. Here we consider another example of the latter by looking for J/Ψ production from a $c\bar{c}$ component in the B meson wavefunction.

To determine the process that is best suited for our purpose, we first consider purely hadronic decays of the B meson with just one $c\bar{c}$ pair in the final state. It is then easy to see from Fig. 1, that hadronic channels with a $c\bar{c}\ X_{u,d,s}$ quark content are also produced at tree level via V_{cb}, with no need for IC. We conclude that these processes are ill-suited to study the effects of IC, which would be swamped by a formidable background. Next, we consider a process with three charm quarks (or antiquarks) in the final state such as $B \to J/\Psi D^{(*)}$ or $B \to J/\Psi D^{(*)} X$. The former has been discussed in Ref. [2] and in the latter the phase space available closes rapidly, suppressing the rates to unobservable levels.



Figure 1: Perturbative contribution to the decay $\overline{B}^0_d \to J/\Psi\, X$. (a) is a standard tree-level graph, (b) is an annihilation graph involving IC in the initial state. An IC exchange graph is also allowed.

This leads us to concentrate in semileptonic modes with a J/Ψ in the final state. These modes have the additional advantage of presenting an extremely clean experimental signature containing three charged leptons. The simplest mode of this type is $B \to J/\Psi\ e^- \overline{\nu}_e X$.[1] Detectors such as CDF operating at hadron machines, or experiments at B factories such as *BABAR* at SLAC should be able to identify the three outgoing charged leptons easily. In fact, CDF will be looking for the mode $B_c \to J/\Psi\ e^- \overline{\nu}_e$ which has the same topology (and which would constitute a background to our process). It would therefore be natural for them to search for $B \to J/\Psi\ e^-\overline{\nu}_e\ X$ simultaneously.

---

[1] For definiteness we concentrate here on channels with a J/Ψ in the final state, but our analysis can also be applied to channels with other $c\overline{c}$ resonances or with $D\overline{D}$ pairs instead.

This paper is organized as follows: in Section 2 we calculate the perturbative QCD contributions to semileptonic processes such as $B^- \to J/\Psi\ e\nu\, X$ and show that they are unobservably small. In Section 3 we propose a modified spectator model for the computation of the IC contributions to these same semileptonic processes, and find that they can be substantially larger, with branching ratios as large as 5×10^{-7}. Finally in Section 4 we estimate the reach of near future experiments for these reactions and show that they could observe the mode $B^- \to J/\Psi\ e\nu\, X$ if the IC component of B mesons is as large as has been speculated recently [4].

2 The perturbative calculation

In this section we compute the perturbative contribution to the decay rate of the process $B^- \to J/\Psi\ e^-\overline{\nu}_e\, X$ from the two lowest order diagrams in Fig. 2. We present calculations of the decay rate in both Non-Relativistic QCD (NRQCD) [7] and in the Color Evaporation Model (CEM)[8]. We also derive the leading order contribution to the total inclusive B-to-$(c\bar{c})$ semileptonic decay rate.

A hadronic B decay depends on the momentum scale of the meson wavefunction ($1/\mathrm{fm} \sim \Lambda_{\mathrm{QCD}}$) and is non-perturbative in principle. Nevertheless, the mass of the heavy quark provides a large momentum scale and to the extent that $m_{J/\psi} \sim 2m_c \gg \Lambda_{\mathrm{QCD}}$ and $m_b - m_{J/\psi} \gg \Lambda_{\mathrm{QCD}}$, the $c\bar{c}$ pair production occurs at short distance and is perturbative. On the other hand, the formation of J/Ψ mesons from the initially compact $c\bar{c}$ pairs takes a long time and is non-perturbative. The quantum interference between the production of the $c\bar{c}$



Figure 2: Graphs describing the perturbative contribution to the decay $B^- \to J/\Psi\, e^-\overline{\nu}\, X$.

pair and the formation of J/Ψ mesons should be suppressed by powers of the ratios of the typical momentum scales of these two processes, such as $\Lambda^2_{\rm QCD}/m^2_{J/\psi}$ and $\Lambda^2_{\rm QCD}/(m_b - m_{J/\psi})^2$. Therefore, if we neglect these power corrections, the decay rate can be factorized as

$$\begin{aligned}\Gamma_{B^-\to J/\psi\, e\bar{\nu}\, X} &= \sum_n \int dq^2 \int d^4Q \left(\frac{d\Gamma_{B\to[c\bar{c}]_n\, e\bar{\nu}\, X}}{d^4Q}\right) \\ &\times\ \delta\left(q^2 + (k_c - k_{\bar{c}})^2\right)\, F_{[c\bar{c}]_n\to J/\psi}(q^2), \qquad (1)\end{aligned}$$

where $\sum_n$ runs over all possible singlet and octet color states, and $Q^\mu = k_c^\mu + k_{\bar{c}}^\mu$ is the total momentum of the produced $c\bar{c}$ pair. In Eq. (1), $q^2 = (2\vec{k}_c)^2$ is the square of the relative momentum between the c and the $\bar{c}$ in their rest frame. Treating the c or $\bar{c}$ approximately as being on their mass-shell, $k_c^2 = k_{\bar{c}}^2 = m_c^2$, $q^2 = (k_c - k_{\bar{c}})^2 = Q^2 - 4\,m_c^2$. The non-perturbative quantity $F_{[c\bar{c}]_n\to J/\psi}(q^2)$ denotes the transition probability for a $[c\bar{c}]$ state with relative momentum squared q^2 to evolve into a physical J/ψ meson. Different choices of $F_{[c\bar{c}]_n\to J/\psi}(q^2)$ lead to different models of J/Ψ formation.

2.1 Nonrelativistic QCD

The NRQCD formalism for quarkonium production exploits the fact that the heavy quark velocity v_c is much less than one, or equivalently that the heavy quark three-momentum is much smaller than its mass ($|\vec{k}_c| \ll m_c$) [7]. By expanding the q^2-dependence of $d\Gamma_{B^- \to [c\bar{c}]_n X}/d^4Q$ around $q^2 = 0$ in Eq. (1), we effectively derive the leading term in the NRQCD formalism:

$$\Gamma(B^- \to J/\Psi\ e^- \overline{\nu}_e\, X) = \sum_n \Gamma(B^- \to [c\bar{c}]_n\, e^- \overline{\nu}_e\, X)\, \langle \mathcal{O}_n^{J/\Psi} \rangle\,. \tag{2}$$

The constants $\langle \mathcal{O}_n^{J/\Psi} \rangle \propto \int dq^2 F_{[c\bar{c}]_n \to J/\psi}(q^2)$ are matrix elements of local operators in coordinate space, often called NRQCD matrix elements. They contain all the non-perturbative information on the J/Ψ formation from the $c\bar{c}$ pair. According to the NRQCD formalism for heavy quarkonium production, the color of the gluon in Fig. 2 is neutralized by non-perturbative soft gluons in the production process, which are not explicitly shown. For the lowest order diagrams in Fig. 2 only a spin-1 and color-octet $c\bar{c}$ state is able to produce a J/Ψ, and the corresponding matrix element is $\langle \mathcal{O}_8^{J/\Psi}(^3S_1) \rangle$ [9].

To the extent that the J/Ψ mass is large, the partonic process in the decay amplitudes of $B^- \to [c\bar{c}]_8\, e^- \overline{\nu}_e\, X$ in Fig. 2 takes place at short distance $\sim 1/m_\Psi$. More precisely, the gluon is off-shell by the J/Ψ mass, and the denominator of the t- or u-channel fermion propagator is at least 5 GeV^2. Therefore, the decay in Fig. 2 can be factorized into a hadronic matrix element for $B \to b\bar{u}$, and a partonic process $b\bar{u} \to [c\bar{c}]_8\, e^- \overline{\nu}_e$. In order to achieve this factorization we formally write the decay amplitudes of $B \to g^\star W^\star$ in Fig. 2 as:

$$M_i^{\alpha\mu} = \int \frac{d^4 p_b}{(2\pi)^4} \operatorname{Tr}\left[\widehat{T}_B(p_B,\ p_b) \cdot \widehat{H}_i^{\alpha\mu}(p_b,\ p_\Psi,\ q)\right], \qquad i = 1,\ 2\ , \tag{3}$$

where $\widehat{T}_B$ and $\widehat{H}$ represent the matrix elements of $B \to b\bar{u}$ and the partonic process $b\bar{u} \to$

$g^\star W^\star$ respectively. The partonic momenta satisfy $p_{\bar{u}} = p_B - p_b$, and the sum of leptonic momenta is $q = q_1 + q_2 = q_e + q_{\overline{\nu}}$.

By decoupling the spinor trace between the matrix element and the partonic process, and boosting p_B to an infinite momentum frame, the decay amplitudes in Eq. (3) can be approximated as ($M_1^{\alpha\mu}$ and $M_2^{\alpha\mu}$ refer to the contributions from the direct and crossed graphs, respectively),

$$M_i^{\alpha\mu} \approx \int dx\, T_B^\sigma(x)\; H_{i\sigma}^{\alpha\mu}(p_b = x p_B,\; p_\Psi,\; q)\;, \tag{4}$$

where the matrix element is defined by

$$T_B^\sigma(x) = \int \frac{p_B^+\, dy^-}{2\pi}\; \exp(i\, x\, p_B^+\, y^-)\; \langle\, 0 \mid \overline{u}(0)\gamma^\sigma\gamma_5\, b(y^-) \mid B^- \rangle\;, \tag{5}$$

and the partonic process H is computed extracting the axial-vector component of the lowest order amplitudes $b\bar{u} \to g^\star W^\star$.

Eq. (5) is given in terms of light-cone coordinates: the ± components of any four-vector k^μ are defined as $k^\pm = (k^0 \pm k^3)/\sqrt{2}$. We also write $p_B^\mu = \left(p_B^+,\; m_B^2/(2p_B^+),\; 0_\perp\right)$ using the boost-invariant light-cone wavefunction language.

Due to the heavy quark mass, the typical virtuality of b and $\bar{u}$ in the B meson should be much smaller than the large momentum scale in H, and therefore we can approximate the parton momenta p_b and $p_{\bar{u}}$ in $H_{i\sigma}^{\alpha\mu}$ as

$$p_b = x\, p_B\;, \qquad p_{\bar{u}} = (1-x)\, p_B\;, \qquad \text{with } x = \frac{m_b}{m_B}\;. \tag{6}$$

This approximation corresponds to dropping power corrections suppressed by the hard momentum scale of the partonic process. Fixing x in $H_{i\sigma}^{\alpha\mu}$ we decouple the x integration in

Eq. (4) to find

$$\int dx\, T^\sigma_B(x) = \langle\, 0 \mid \overline{u}(0)\gamma^\sigma\gamma_5\, b(0) \mid B^- \rangle \;\equiv i f_B\, p^\sigma_B\,. \tag{7}$$

For numerical estimates we use $f_B = 0.177$ GeV [10, 11][2]. Substituting Eq. (7) into Eq. (4) we obtain a factorized expression for the decay amplitude of $B \to g^\star W^\star$,

$$M^{\alpha\mu}_i \approx (i f_B\, p^\sigma_B)\;\; H^{\alpha\mu}_{i\sigma}\left(p_b = x p_B,\; p_\Psi,\; q\right)\Big|_{x=\frac{m_b}{m_B}}\,. \tag{8}$$

The amplitudes in Eq. (3) can thus be written as

$$\begin{aligned} M^{\alpha\mu}_1 &= f_B\,\frac{g_S g_W}{2\sqrt{2}}\frac{1}{(p_\Psi - p_{\bar{u}})^2 - m^2_{\bar{u}}}\,V_{ub}\left[-i\,\varepsilon^{\gamma\delta\alpha\mu} p_{B\gamma}(p_\Psi - p_{\bar{u}})_\delta\right. \\ &+ \left. p^\mu_B (p_\Psi - p_{\bar{u}})^\alpha + p^\alpha_B (p_\Psi - p_{\bar{u}})^\mu - p_B\cdot(p_\Psi - p_{\bar{u}})\, g^{\alpha\mu}\right]\;, \end{aligned} \tag{9}$$

and (for the crossed diagram),

$$\begin{aligned} M^{\alpha\mu}_2 &= f_B\,\frac{g_S g_W}{2\sqrt{2}}\frac{1}{(q - p_{\bar{u}})^2 - m^2_b}\,V_{ub}\left[-i\,\varepsilon^{\gamma\delta\alpha\mu} p_{B\gamma}(p_{\bar{u}} - q)_\delta\right. \\ &+ \left. p^\mu_B (-p_{\bar{u}} + q)^\alpha + p^\alpha_B (-p_{\bar{u}} + q)^\mu - p_B\cdot(-p_{\bar{u}} + q)\, g^{\alpha\mu}\right]\;. \end{aligned} \tag{10}$$

We denote the strong and weak coupling constants by g_S and g_W respectively.

The $W^\star$ coupling to the lepton current is evaluated in the usual way. After squaring this contributes a factor

$$L^{\alpha\beta} = g^2_W\,\left(\frac{1}{q^2 - m^2_W}\right)^2\,\left[i\varepsilon^{\gamma\delta\alpha\beta} q_{1\gamma} q_{2\delta} + q^\beta_1 q^\alpha_2 + q^\alpha_1 q^\beta_2 - q_1\cdot q_2\, g^{\alpha\beta}\right] \tag{11}$$

---

[2] $f_B = 0.177$ GeV corresponds to the usual annihilation of a $b\bar{u}$ pair in the color singlet configuration. In our case the $b\bar{u}$ pair is in a color octet configuration and therefore we expect that our f_B is actually smaller than 0.177. Our results for the perturbative production of J/Ψ in semileptonic B decays can thus be considered to be an upper bound.

to the decay rate.

Our remaining task is to parametrize the production of a $J/\Psi(^3S_1)$ state from the $c\overline{c}$ pair. We do this with projector techniques and compute the overall color factor below (in Eq. (15) [9]). We write

$$J_{\mu\nu}(p_\Psi^2) = \sum_\lambda \left(\frac{-i}{p_\Psi^2}\right) \mathrm{Tr}[(-ig_S\gamma_\mu)\,\Pi_1^\rho]\;\varepsilon_\rho^{(\lambda)}(p_\Psi)\cdot\left(\frac{i}{p_\Psi^2}\right)\mathrm{Tr}[(ig_S\gamma_\nu)\,\Pi_1^{*\sigma}]\;\varepsilon_\sigma^{*(\lambda)}(p_\Psi)\;, \tag{12}$$

with the projector defined as [9]

$$\Pi_1^\lambda = \frac{1}{\sqrt{m_\Psi^3}}\frac{1}{4}\left(\not{p}_\Psi - m_\Psi\right)\gamma^\lambda\left(\not{p}_\Psi + m_\Psi\right)\;. \tag{13}$$

The total contribution to the decay rate from the $c\overline{c}$ line is then

$$J_{\mu\nu}(p_\Psi^2) = 4\,g_S^2\left(\frac{1}{p_\Psi^2}\right)^2\frac{1}{m_\Psi}\left[p_\mu^\Psi p_\nu^\Psi - m_\Psi^2 g_{\mu\nu}\right]\;, \tag{14}$$

and the overall color factor is

$$\frac{1}{N_c^2-1}\sum_{ABC}\mathrm{Tr}[T^A C_8^B]\;\mathrm{Tr}[T^A C_8^C] = \frac{1}{2}\;,\qquad C_8^A = \sqrt{2}\,T^A\;, \tag{15}$$

where T^A, A = 1...8, are the generators of the color group.

As shown in Eq. (2), the rate of the process $B^- \to J/\Psi\, e^-\overline{\nu}_e\, X$ includes the NRQCD matrix element $\langle\mathcal{O}_8^{J/\Psi}(^3S_1)\rangle$ defined in [9, 12, 13], which represents the probability for an almost point-like $c\bar{c}$ pair in a color-octet $^3S_1$ state to bind and form a J/Ψ. The matrix element is nonperturbative and has to be extracted from experimental data, and its numerical value is [9, 12]:

$$\langle\mathcal{O}_8^{J/\Psi}(^3S_1)\rangle = 1.19\times10^{-2}\ \mathrm{GeV}^3\;. \tag{16}$$

As a final step to obtain the decay rate we need to divide by the number of color and polarization states of the $c\bar{c}(^3S_1)$ produced, $N_{col}N_{pol} = 24$, as explained in Ref. [9].

Collecting our results, we obtain for the matrix element squared

$$|\mathcal{M}|^2 = (M_1^{\alpha\mu} + M_2^{\alpha\mu})^\star \left(M_1^{\beta\nu} + M_2^{\beta\nu}\right) L_{\alpha\beta}\, J_{\mu\nu}(p_\Psi^2)\, \frac{1}{2}\, \frac{\langle \mathcal{O}_8^{J/\Psi}(^3S_1)\rangle}{N_{col}N_{pol}}\ , \tag{17}$$

which includes the color factor 1/2 obtained in Eq. (15). In our numerical results we use the boson masses [14] $m_B = 5.279$ GeV, $m_\Psi = 3.097$ GeV, and the current quark mass $m_b = 4.9$ GeV, together with $|V_{ub}| = 0.0035$ [15], $\alpha_S(2m_c) = 0.266$ [16] and $\tau_{B^-} = 1.653 \times 10^{-12}$ sec [14]. We find a branching ratio:

$$\mathrm{BR}(B^- \to J/\Psi\ e^-\overline{\nu}_e\, X) = 9 \times 10^{-11}\ . \tag{18}$$

The process in Fig. 2 which we have calculated is the simplest one giving rise to $B^- \to J/\Psi\ e^-\overline{\nu}_e\ X$. However, it suffers from a suppression factor f_B associated with the $b\bar{u}$ annihilation. For this reason it is possible that a process involving more quarks but no annihilation might actually be larger. To investigate this possibility we now estimate the rate of the process $B \to J/\Psi\ e^-\overline{\nu}_e\ X_{u,c}$ as in Fig. 3.

We adopt once more the NRQCD factorization form of Eq. (2) to calculate separately the matrix elements of $b \to q\ g^\star e^-\overline{\nu}_e$ $(q = u, c)$, and for the conversion of the $c\bar{c}$ pair into a J/Ψ. We denote by $H_1^{\alpha\mu}$ and $H_2^{\alpha\mu}$ the amplitudes of the partonic processes where the gluon is emitted from the b and q lines respectively. The squared partonic matrix element is then

$$M^{\alpha\mu\beta\nu} = \mathrm{Tr}\left[(H_1^{\alpha\mu} + H_2^{\alpha\mu})(H_1^{\beta\nu} + H_2^{\beta\nu})^\dagger\right], \tag{19}$$



Figure 3: Spectator graph analogous to Fig. 2 for $B \to J/\Psi\ e^- \overline{\nu}\ X_{u,c}$. The gluon propagator may also be attached to the outgoing quark u or c.

with

$$\begin{aligned} H_1^{\alpha\mu} &= \frac{g_S g_W}{2\sqrt{2}} \frac{1}{(p_b - p_\Psi)^2 - m_b^2} V_{uq} \overline{u}_q(p_q) \gamma^\alpha (1-\gamma_5)(\not{p}_b - \not{p}_\Psi) \gamma^\mu u_b(p_b)\ , \\ H_2^{\alpha\mu} &= \frac{g_S g_W}{2\sqrt{2}} \frac{1}{(p_b - q)^2 - m_q^2} V_{uq} \overline{u}_q(p_q) \gamma^\mu (\not{p}_b - \not{q}) \gamma^\alpha (1-\gamma_5) u_b(p_b)\ . \end{aligned} \tag{20}$$

The leptonic and hadronic $c\overline{c}$ parts and the NRQCD matrix element are the same as for $B^- \to J/\Psi\ e^- \overline{\nu}_e X$, and have been given in Eqs. (11), (14), and (16), respectively. The overall color factor is different,

$$\frac{1}{N_c} \sum_{ABC} \mathrm{Tr}[T^A C_8^C]\ \mathrm{Tr}[T^B C_8^C]\ \sum_{ij} \left(T^A\right)_{ij} \left(T^B\right)_{ij} = \frac{2}{3}\ , \qquad C_8^A = \sqrt{2}\, T^A\ . \tag{21}$$

All these factors are combined into the squared matrix element,

$$|\mathcal{M}|^2 = M^{\alpha\mu\beta\nu} L_{\alpha\beta}\ J_{\mu\nu}(p_\Psi^2)\ \frac{2}{3}\ \frac{\langle \mathcal{O}_8^{J/\Psi}(^3S_1)\rangle}{N_{col} N_{pol}}\ . \tag{22}$$

Using the masses already defined plus [14] $m_c = 1.25$ GeV and $|V_{bc}| = 0.041$, and a b quark lifetime $\tau_b = 1.6 \times 10^{-12}$ sec we find:

$$\mathrm{BR}(B \to J/\Psi\ e^-\overline{\nu}_e\ X_q) = \begin{cases} 1 \times 10^{-12} & \text{for } q = c\ , \\ 4 \times 10^{-12} & \text{for } q = u\ . \end{cases} \tag{23}$$

These results show that the diagrams in Fig. 2 generate the leading perturbative contribution to the process $B \to J/\Psi\ e^-\overline{\nu}_e\ X_q$, which is unobservably small.

2.2 Total inclusive B-to-$c\bar{c}$ semileptonic decay

The inclusive semileptonic rate for $B^- \to c\bar{c}\ e^-\overline{\nu}_e\, X$ is obtained from the diagrams in Fig. 2 with some simple replacements. The calculation proceeds as in Section 2.1 up until Eq. (11). After that we simply couple the $g^\star$ to a $c\bar{c}$ line in the usual way, obtaining

$$J'_{\mu\nu}(p_1, p_2) = 4\, g_S^2 \left(\frac{1}{p^2}\right)^2 \Big[p_{1\mu}p_{2\nu} + p_{2\mu}p_{1\nu} - (p^2/2)g_{\mu\nu}\Big]\ , \tag{24}$$

where p_1 and p_2 are the momenta of c and $\bar{c}$, respectively, and $p = p_1 + p_2$. The overall color factor now takes into account both the color octet and the color singlet configurations of the incoming quarks b and $\bar{u}$, and can be calculated to be 2/9. The squared matrix element becomes

$$|\mathcal{M}|^2 = (M_1^{\alpha\mu} + M_2^{\alpha\mu})^\star \left(M_1^{\beta\nu} + M_2^{\beta\nu}\right) L_{\alpha\beta}\ J'_{\mu\nu}(p_1, p_2)\, \frac{2}{9}\ , \tag{25}$$

where $M_i^{\alpha\mu}$ and $L_{\alpha\beta}$ are the same quantities already used in Eq. (17).

We use the masses and CKM parameters defined previously, but we require the invariant mass of the $c\bar{c}$ pair to be larger than 3 GeV in order to produce a physical $c\bar{c}$ meson.

We obtain a branching ratio

$$\mathrm{BR}(B^- \to c\bar{c}\, e^-\overline{\nu}_e\, X) = 4.2 \times 10^{-10}. \qquad (26)$$

The differential branching ratio is plotted in Fig. 4 as a function of the $c\bar{c}$ pair invariant mass. From Fig. 4 we can see that about 90% of the time the $c\bar{c}$ pair will become a charmonium resonance and only about 10% of the time it will lead to $D\bar{D}$ production.



Figure 4: The differential branching ratio of $B^- \to c\bar{c}\, e^-\overline{\nu}\, X$ is plotted vs. the invariant mass $m_{c\bar{c}}$ of the $c\bar{c}$ pair. The vertical dashed line shows the energy threshold for $D\bar{D}$ production.

2.3 Color evaporation model

According to the Color Evaporation model of J/Ψ production, all the $c\bar{c}$ pairs with invariant mass below open-charm threshold should become charmonia by radiating soft gluons to neutralize the color and other quantum numbers [8]. The formation process from the $c\bar{c}$ pairs to J/Ψ mesons is assumed to be independent of the color and invariant mass of the pairs. This is equivalent to choosing

$$F_{[c\bar{c}]_n \to J/\Psi}(q^2) = f_{J/\Psi}\, \theta(4m_D^2 - 4m_c^2 - q^2)\,, \tag{27}$$

in Eq. (1). Here $4m_D^2$ represents the open-charm threshold, and $f_{J/\Psi}$ is a universal constant encoding the relative probability for a $c\bar{c}$ pair to become a J/Ψ instead of other charmonium states, and needs to be fixed by data. Substituting Eq. (27) into Eq. (1), we derive the formula for the decay rate in the Color Evaporation model,

$$\Gamma_{B^- \to J/\psi\; e\bar{\nu} X} = f_{J/\Psi} \int_{2m_c}^{2m_D} dm_{c\bar{c}} \left[\frac{d\Gamma_{B^- \to c\bar{c}\; e\bar{\nu} X}(m_{c\bar{c}})}{dm_{c\bar{c}}} \right]\,, \tag{28}$$

where $m_{c\bar{c}}$ is the invariant mass of the $c\bar{c}$ pair. Using the results of our previous section shown in Fig. 4, this becomes

$$\mathrm{BR}(B^- \to J/\psi\; e\bar{\nu}\, X) = f_{J/\Psi}\; 3.8 \times 10^{-10}\,. \tag{29}$$

Comparing Eq. (29) with Eq. (18) we see that the CEM model result would agree with the NRQCD result if $f_{J/\Psi} \sim 1/4.3$. This fits very well with the expectation that $1/f_{J/\Psi} \sim 2$ to 7 in Ref. [8].

Our results in Eqs. (18) and (23) show that the perturbative QCD contributions to the processes $B^- \to J/\Psi\; e^-\overline{\nu}_e\, X$ and $B \to J/\Psi\; e^-\overline{\nu}_e\; X_{u,c}$ are too small to be observed by near-future experiments. Qualitatively, these results are so small because they involve two off-shell propagators as well as suppressions from powers of α_S and from the octet matrix element $\langle \mathcal{O}_8^{J/\Psi}(^3S_1) \rangle$. This leaves open the possibility of significant enhancements due to IC which we explore in the following section.

3 The intrinsic charm contribution

The mechanism by which IC in a B meson leads to the decay $B^- \to J/\Psi\ e^-\overline{\nu}_e X$ is different from the perturbative case. The two (anti)charm quarks are now part of the initial wavefunction as shown in Fig. 5. In this picture a fluctuation in an ordinary B^- meson produces a nearly on-shell $c\bar{c}$ pair which is part of the initial wavefunction. There is a small probability for this fluctuation to occur [1], but once it happens, it is possible for the off-shell $b\bar{u}$ quarks to annihilate leaving a J/Ψ behind. In this section we attempt to estimate the rate of this process in the following way. We start from a B^- meson wavefunction in the higher Fock state $|b\bar{u}c\bar{c}\rangle$ with a certain momentum distribution as proposed in the IC picture of Brodsky *et al.* [1]. We then estimate the annihilation rate of the $b\bar{u}$ components treating the $c\bar{c}$ as spectators. This annihilation is negligible unless the $b\bar{u}$ system is in a $J = 1$ state, and in this case we think of it as an off-shell $B^\star$. As we will see, the invariant mass of the remaining $c\bar{c}$ pair is dominantly in the charmonium resonance region, so we assume that a certain fraction produces a J/Ψ as in the CEM model discussed in the previous section.



Figure 5: IC contribution to the decay $B^- \to J/\Psi\ e^-\overline{\nu}\,X$.

The momentum fraction distribution of the $|b\bar{u}c\bar{c}\rangle$ light-cone Fock component of B^- implies that IC carries large momentum fractions, unlike the small x of extrinsic quarks from gluon splitting. It is given by [1]:

$$\frac{dP_{IC}^{B}}{dx_1\cdots dx_4} = N_4 \frac{\delta(1-\sum_{i=1}^{4} x_i)}{(m_B^2 - \widehat{m}_{\bar{u}}^2/x_{\bar{u}} - \widehat{m}_b^2/x_b - \widehat{m}_c^2/x_c - \widehat{m}_{\bar{c}}^2/x_{\bar{c}})^2} \; . \tag{30}$$

In Eq. (30) the x_i are the light-cone momentum fractions $(k^0 + k^3)_i/(p_B^0 + p_B^3)$, and $\widehat{m}_i = (m_i^2 + \langle k_\perp^2 \rangle_i)^{1/2}$ are the transverse masses of each constituent as in Refs. [1]. In our calculation we use the numerical values $\widehat{m}_b = 5$ GeV, $\widehat{m}_{u,d} = 0.45$ GeV and $\widehat{m}_c = 1.8$ GeV [1, 17]. Integrating Eq. (30) over all possible momenum fractions gives the probability R_{IC} of having the B meson in an IC configuration. In terms of this R_{IC} we obtain the four-particle normalization factor $N_4 = R_{IC} \times 10^5$ GeV^4. It has been suggested in Ref. [4] that IC effects in B mesons may be four times larger than in the proton, where it is estimated that they can occur at the 1% level [6]. In particular, this means that R_{IC} could be as large as 0.04 according to Ref. [4].

We are interested in a process in which the $c\bar{c}$ is in the particular configuration ${}^3S_1$ color singlet state. There is, of course, a model dependent combinatorial factor associated with this. Because we lack a detailed dynamical model for IC, we will instead resort to a phase-space model similar in spirit to the CEM discussed in the previous section.

We begin our estimate of the amplitude by treating the charm (anti)quarks as spectators. The wavefunction is then the product of a $(c\overline{c})$ state that generates the J/Ψ and a virtual $(b\overline{u})$ state with the quantum numbers of a B^{*-} (other configurations are helicity suppressed) which decays into leptons. This virtual $(b\overline{u})$ pair possesses an invariant mass squared $m_{B^\star}^2 = q^2 = (x_b + x_{\bar{u}})^2 m_B^2$. The amplitude squared reduces to the product of a leptonic part $L_{\mu\nu}$ which is the same as that in Eq. (11) and the hadronic b–u line contribution. Combining them we find:

$$|\mathcal{M}|^2 = -f_{B^\star}^2 m_{B^\star}^2 \frac{g_W^2}{8} |V_{ub}|^2 g^{\mu\nu} L_{\mu\nu} \ = 4 f_{B^\star}^2 q^4 G_F^2 \ |V_{ub}|^2 \ . \tag{31}$$

It is evident from this result that we treat the $b\overline{u}$ annihilation as the leptonic decay of an off-shell $B^\star$. For our numerical estimates we take $f_{B^\star} = f_B$.

It remains to integrate Eq. (31) over the phase space available to the lepton pair (determined by $m_{B^\star}^2$) and to convolute this result with the momentum distribution in the B meson wavefunction, Eq. (30). In this convolution we impose kinematic constraints consistent with "liberating" the IC pair which becomes a charmonium resonance or a $D\bar{D}$ pair.

Numerically, all this results in the differential decay rate shown in Fig. 6, where we have used $R_{IC} = 0.04$ for illustration. The integrated rate leads to the inclusive branching ratio,

$$\mathrm{BR}_{IC}(B^- \to (c\bar{c})\ e^-\overline{\nu}_e\, X) = 5 \times 10^{-7} \times \frac{R_{IC}}{0.04} \ . \tag{32}$$

If instead we only integrate over the charmonium resonance region, we obtain the estimate,

$$\mathrm{BR}_{IC}(B^- \to J/\Psi\ e^-\overline{\nu}_e\, X) = 4 \times 10^{-7} \times \frac{R_{IC}}{0.04} \ . \tag{33}$$

Only a fraction of this result corresponds to J/Ψ, and if we use our perturbative calculations as guidance, this fraction could be of order $f_{J/\Psi} \sim 1/4.3$. However, as a signal for IC any charmonium resonance that can be reconstructed (or for that matter any $D\bar{D}$ channel) will serve. Integrating the distribution in Fig. 6 above open-charm threshold leads to the



Figure 6: The IC contribution to the differential branching ratio of $B^- \to c\bar{c}\, e^-\overline{\nu}\, X$ is plotted vs. the invariant mass $m_{c\bar{c}}$ of the $c\bar{c}$ pair. The vertical dashed line shows the energy threshold for $D\bar{D}$ production.

estimate $\mathrm{BR}_{IC}(B^- \to D\bar{D}\ e^-\overline{\nu}_e\, X) = 6 \times 10^{-8}\ R_{IC}/0.04$. If the fraction of IC in the B meson can really be as large as $R_{IC} = 0.04$, we see that semileptonic B decay constitutes a powerful probe of these ideas. Our result, Eq. (33), is three orders of magnitude larger than conventional, perturbative, mechanisms for producing charm pairs in semileptonic B meson decay. This large enhancement can be understood qualitatively because the PQCD mechanism is suppressed with respect to the IC mechanism by two powers of α_S, two propagators with an m_Ψ mass scale, and the matrix element for J/Ψ formation from a color octet $c\bar{c}$. At the same time the only relative suppression in the IC mechanism is the probability of finding the B wave-function in the IC configuration, R_{IC}. Roughly, the ratio of PQCD to

IC induced rates goes as $\alpha_S^2(\langle \mathcal{O}_8^{J/\Psi}(^3S_1)\rangle/m_\Psi^3)/R_{IC} \sim 10^{-3}$.

We can perform a similar calculation for the decay $B \to J/\Psi\ e^-\overline{\nu}_e\ X_{u,c}$, depicted in Fig. 7. This process has at least one more particle in the final state and we might expect it to be suppressed by phase space with respect to $B \to J/\Psi\ e^-\overline{\nu}_e\, X$. However, it does not suffer from the suppression factor associated with the weak annihilation of the $b\bar{u}$ pair and for this reason we estimate it here. Our treatment of IC is similar to the previous channel. We calculate the semileptonic decay of an off-shell b-quark into three particles, $b \to e^-\overline{\nu}_e$ (u or c) and treat the remainder three quarks in the B wavefunction as spectators.

As in our previous example, we picture a nearly on-shell $c\bar{c}$ pair arising as a fluctuation in the B-meson wavefunction. The semileptonic decay of the b-quark is standard, the only



Figure 7: IC contribution to the three-body spectator semileptonic decays $b \to J/\Psi\ e^-\overline{\nu}_e$ $\mathrm{X}_{u,c}$.

difference here being that the b-quark is off-shell and carries a fraction x_b of the B meson momentum. The partonic decay rate is thus (in terms of $m_b^\star = x_b m_B$):

$$\Gamma(b \to q\, e^- \overline{\nu}_e) \approx \frac{G_F^2 |V_{bq}|^2 m_b^{\star 5}}{192\pi^3} f\left(\frac{m_q}{m_b^\star}\right) \ , \quad \mathrm{q} = \mathrm{u}, \mathrm{c}\ , \tag{34}$$

where for the case of a V_{bc} transition we also include the usual kinematic factor

$$f(x) = 1 - 8x^2 + 8x^6 - x^8 - 24x^4 \ln x. \tag{35}$$

This factor is approximately equal to 1 for the V_{bu} transition. This result is now folded with the momentum distribution in the B wavefunction, Eq. (30), with kinematic constraints to once again “liberate” the IC pair. In spite of the CKM suppression V_{ub}/V_{cb}, the V_{ub} mode can be as large or larger than the V_{cb} mode because there is a very large phase-space suppression producing three charm quarks. This suppression is so strong that the result is very sensitive to the value of m_c. Using $m_c = 1.25$ GeV and a b quark lifetime $\tau_b = 1.6 \times 10^{-12}$ sec, we estimate

$$\mathrm{BR}_{IC}(B \to J/\Psi\ e\nu_e\ X_q) \approx \begin{cases} 1 \times 10^{-7} \times R_{IC}/0.04 & \text{for } q = c\ , \\ 4 \times 10^{-8} \times R_{IC}/0.04 & \text{for } q = u\ . \end{cases} \tag{36}$$

If instead we use $m_c = 1.6$ GeV we obtain $\mathrm{BR}_{IC}(b \to J/\Psi\ e^- \overline{\nu}_e\ X_c) = 4 \times 10^{-9} \times R_{IC}/0.04$. From this we conclude that the V_{ub} mode is a small correction to our previous process so that $B^- \to J/\Psi\ e^- \bar{\nu}\ X_u$ is dominated by the annihilation mode $B^- \to J/\Psi\ e^- \bar{\nu}\, X$. We also conclude that the channel with three charm quarks in the final state is smaller in spite of the CKM advantage it enjoys.

4 Discussion and conclusions

Up to now we have discussed the production of a $c\bar{c}$ state like J/Ψ in a semileptonic B^- decay. We have shown that the perturbative QCD contributions to this process are extremely small, giving rise to branching ratios at the 10^{-10} level. Similar conclusions apply to the perturbative production of other charmonium resonances or $D\bar{D}$ pairs. We have also shown that there can be significantly larger contributions from IC components of the B-meson wavefunction.

It is simple to understand why the IC contribution to our processes is much larger than the perturbative contribution. The perturbative contribution suffers from several suppression factors associated with the exchange of a perturbative gluon and the color rearrangements that are needed to convert two color octet $q\bar{q}$ pairs into color singlet mesons. The IC contribution bypasses all this because all necessary color exchanges and rearrangements take place in a nonperturbative way inside the B-meson wavefunction. The only suppression for this mechanism is then the small probability of finding the B meson in the IC configuration.

Adopting a simple spectator model and accepting recent suggestions in the literature as to the possible size of the IC component, we have estimated that $B^- \to (c\bar{c})\ e^-\bar{\nu}X$ could have a branching ratio as large as 5×10^{-7}. We now show that at this level, the process $B^- \to (c\bar{c})\ e^-\bar{\nu}X$ could be observed in the near future.

We start by estimating the sensitivity of CDF to the process $B \to J/\Psi\ e\nu X$ in Run II by comparing it to $B_c \to J/\Psi\ l\nu$, which they expect to see and which has the same final state. Assuming 2 fb^{-1} and 15 fb^{-1} respectively for the integrated luminosities during Run IIa and Run IIa+Run IIb, and using [18]

$$\frac{\text{BR}(b \to B_c)}{\text{BR}(b \to B^-)} \times \frac{\text{BR}(B_c \to J/\Psi\ l\nu)}{\text{BR}(B^- \to J/\Psi\ K^-)} = 0.13 \pm 0.06\ , \tag{37}$$

leads to the estimate [3]

$$N(B_c \to J/\Psi\, l\nu) \sim 800\, \frac{\int \mathcal{L}}{2\ \mathrm{fb}^{-1}} = 6000\, \frac{\int \mathcal{L}}{15\ \mathrm{fb}^{-1}}\ . \tag{38}$$

From this estimate, with $l = \mu$, e, the number of events for $B^- \to J/\Psi\, l\nu\, X$ is given by

$$N(B \to J/\Psi\, l\nu\, X) \sim 5 \times \frac{\int \mathcal{L}}{15\ \mathrm{fb}^{-1}} \times \frac{\mathrm{BR}_{IC}(B \to J/\Psi\, l\nu\, X)}{10^{-7}}\ . \tag{39}$$

Therefore, CDF could be able to use $B \to J/\Psi\ l\nu\, X$ to test IC in the upcoming Run II. Notice that the open-charm channels such as $B \to D^+D^-\ l\nu\, X$ could also be used. Our modeling of IC predicts a smaller branching ratio, of order 6×10^{-8}. However, the identification of charged D mesons could be up to 10 times more efficient than that of J/Ψ since the latter is reconstructed from its leptonic decay which has a small branching ratio, $B(J/\Psi \to \mu^+\mu^- + e^+e^-) \approx 0.12$ [14].

A similar projection for *BABAR* can be based on the decay $B^0 \to D^\star\ l\nu$. Given the 7517 ± 104 reconstructed events for $B^0 \to D^\star\ l\nu$ with a total integrated luminosity of 9.7 fb^{-1} [19], and a projected total luminosity of $\sim 542\ \mathrm{fb}^{-1}$ for year 2006, we have:

$$N(B^0 \to D^\star\ l\nu) \sim 4.2 \times 10^5\, \frac{\int \mathcal{L}}{542\ \mathrm{fb}^{-1}}\ , \tag{40}$$

and from this we can estimate

$$\begin{aligned} N(B \to J/\Psi\ e\nu\, X) \ &\sim\ \mathrm{BR}(J/\Psi \to\ ee + \mu\mu) \times N(B^0 \to D^\star\ l\nu) \times \frac{\mathrm{BR}_{IC}(B \to J/\Psi\ e\nu\, X)}{\mathrm{BR}(B^0 \to D^\star\ l\nu)} \\ &\sim\ 1 \times \frac{\mathrm{BR}_{IC}(B \to J/\Psi\ e\nu\, X)}{10^{-6}}\ . \end{aligned} \tag{41}$$

---

[3] We are grateful to T. LeCompte and M. Tanaka for providing us with this estimate.

We have included a factor $\mathrm{BR}(J/\Psi \rightarrow \ ee + \mu\mu)$ to roughly account for the lower reconstructing efficiency of J/Ψ compared to $D^\star$. From these numbers we conclude that $\mathrm{BR}_{IC}(B \rightarrow J/\Psi\ e\nu\, X)$ could be probed at the 10^{-7} level at CDF and at the 10^{-6} level at *BABAR*. Similar considerations can be applied to Belle.

Near future experiments could use semileptonic B-meson decays to probe the ideas of IC at an interesting level. Their study would complement the suggestion of Ref. [2] and can place limits on the IC content of B mesons.

Acknowledgments

This work was supported in part by the U.S. Department of Energy under Contract Numbers DE-FG02-01ER41155 and DE-FG02-87ER40371. We also thank James Cochran, Thomas LeCompte, Masa Tanaka, Fabio Maltoni and Aida El-Khadra for discussions.

References

[1] S. J. Brodsky, P. Hoyer, C. Peterson and N. Sakai, Phys. Lett. B **93**, 451 (1980); S. J. Brodsky, C. Peterson and N. Sakai, Phys. Rev. D **23**, 2745 (1981).

[2] C.-H. V. Chang and W.-S. Hou, Phys. Rev. D **64**, 071501 (2001).

[3] A. A. Petrov, Phys. Rev. D **58**, 054004 (1998).

[4] S. J. Brodsky and S. Gardner, Phys. Rev. D **65**, 054016 (2002).

[5] S. J. Brodsky and M. Karliner, Phys. Rev. Lett. **78**, 4682 (1997).

[6] B. W. Harris, J. Smith, and R. Vogt, Nucl. Phys. **B461**, 181 (1996).

[7] G. T. Bodwin, E. Braaten, and G. P. Lepage, Phys. Rev. D **51**, 1125 (1995); Erratum *ibid.* **55**, 5853 (1997).

[8] H. Fritzsch, Phys. Lett. **B67**, 217 (1977); M. Glück, J.F. Owens, and E. Reya, Phys. Rev. **D17**, 2324 (1978).

[9] M. Beneke, F. Maltoni, and I. Z. Rothstein, Phys. Rev. D **59**, 054003 (1999); F. Maltoni, M. L. Mangano, and A. Petrelli, Nucl. Phys. **B519**, 361 (1998).

[10] C. W. Bernard, Nucl. Phys. (Proc. Suppl.) **94**, 159 (2001).

[11] C. Bernard *et al.*, Phys. Rev. D **65**, 014510 (2002).

[12] M. Beneke, Lectures given at the *24th Annual SLAC Summer Institute on Particle Physics: The Strong Interaction, From Hadrons to Protons* (SSI 96), Stanford, CA, in “Stanford 1996, The strong interaction, from hadrons to partons”, 549-574.

[13] E. Braaten, S. Fleming, and T. C. Yuan, Annu. Rev. Nucl. Part. Sci. **46**, 197 (1996).

[14] Particle Data Group, D. E. Groom *et al.*, Eur. Phys. J. C **15**, 1 (2000).

[15] A. J. Buras, Introductory lecture given at *KAON2001: International Conference on CP Violation*, Pisa, Italy, 12-17 Jun 2001, hep-ph/0109197.

[16] G. Eilam, M. Ladisa, and Y.-D. Yang, Phys. Rev. D **65**, 037504 (2002).

[17] R. Vogt, S. J. Brodsky, and P. Hoyer, Nucl. Phys. **B360**, 67 (1991); *ibid.* **383**, 643 (1992).

[18] CDF Collab., K. Anikeev *et al.*, preprint FERMILAB-Pub-01/197, December 2001, hep-ph/0201071.

[19] *BABAR* Collab., B. Aubert *et al.*, SLAC-PUB-8530, BABAR-CONF-00-08, Aug 2000, Talk given at the *30th International Conference on High-Energy Physics*, Osaka, Japan, 27 Jul - 2 Aug 2000, hep-ex/0008052.